



Specification List

Autoimmune Diagnostics

General Information

The specifications list contains the range of analysis provided by the clinical immunological laboratory and informs you about the methods, sample types and volumes used, the frequency of analysis, reference values and, if necessary, preanalytical aspects. The specifications list is available on the internet at www.labor-stoecker.de.

Duration:

The samples are registered upon receipt in the laboratory. Preferably, analysis is performed or started on the same day to ensure that the report can be sent to the customer generally one day after receipt of the sample at the latest. The time period may be longer in individual cases if the analysis is not carried out daily or retesting of the sample or performing of a dilution series is required.

Reference range:

The reference values given in the specifications list are in accordance with the manufacturer's recommendations for the test system used and with current scientific and technological knowledge. However, reference ranges may change due to new scientific findings and technical developments in analytical techniques. Interpretation of results is always based on the reference ranges and additional remarks which are given on the report.

Measurement deviations:

The clinical immunological laboratory performs regular checks and maintenance to minimise variations and method-related deviations in the measurement results. Due to existing measurement deviations, borderline and weak-positive results or results that lie near to the reference value or other medical decision limits should be interpreted with care. Information about measurement deviations in individual quantitative analyses can be provided upon request. However, most analyses that are performed by the clinical immunological laboratory (antibody diagnostics) are considered qualitative.

Comparability of test results:

Results from different methods (e.g. immunofluorescence, ELISA, blot techniques) for detecting the same antibodies in a sample may differ. This can be due to the use of different antigens or to varying immobilization of the same antigen to different carrier materials, which can influence the binding of the antibody to the antigen through conformational changes in the antigen or through steric inhibition.

Laboratory developed test (LDT):

The laboratory tests are usually CE marked. Tests that are not CE marked but validated by the laboratory are identified in the specification list as laboratory developed tests (LDT) and are not covered by the accreditation of the clinical immunological laboratory.



21-Hydroxylase IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Acetylcholine-receptor IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA, IFT
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Acetylcholine receptors

sample types: serum
sample volume: 1,5 ml
method: RIA
frequency: biweekly
reference value: see result report
as at: 12/21

Aggrin IgG


sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is a LDT and not covered by the accreditation of our laboratory.
as at: 06/22

Actin (F-Actin) IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Actin IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21



Adrenal cortex IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

AK5 IgG

sample types: serum, EDTA, heparin, citrat plasma
alternative sample types: CSF
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is a LDT and not covered by the accreditation of our laboratory.
as at: 06/23

Alveolar basement membrane IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is not covered by the accreditation of our laboratory.
as at: 05/22

AMA IgG


sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

AMA M2-3E IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

AMA Profil IgGM (M2-3e, M2, M4, M9)

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: EUROLINE
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21



AMA-M7 (myocard specific) IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Amphiphysin IgG

sample types: serum, EDTA, heparin, citrat plasma
alternative sample types: CSF (LDT, not accredited)
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 05/22

ANA IgG (autoantibodies against nuclei) IF global test

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

ANA Profile 23 IgG (nukleosomes, dsDNA, histones, SS-A, Ro-52, SS-B, nRNP/Sm, Sm, Mi alpha, Mi beta, Ku, CENP A, CENP B, Sp100, PML, Scl-70, Pm-Scl100, Pm-Scl75, RP11, RP155, gp210, PCNA, DFS70)


sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: EUROLINE
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

ANCA (autoantibodies against granulocyte cytoplasm)

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

ANCA IgG (MPO, PR3, GBM)

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: EUROLINE
frequency: daily (mo-fri)
reference value: see result report
as at: 11/24



ANCA Profile IgG (PR3, MPO, Elastase, Cath.G, BPI, Lactoferrin)

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

ANNA-3 IgG

sample types: serum, EDTA, heparin, citrat plasma
alternative sample types: CSF (LDT, not accredited)
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 05/22

AP3B2 IgG

sample types: serum, EDTA, heparin, citrat plasma
alternative sample types: CSF
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is a LDT and not covered by the accreditation of our laboratory.
as at: 06/22

Aquaporin-4 IgG


sample types: serum, EDTA, heparin, citrat plasma
alternative sample types: CSF
sample volume: 2,0 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

ASCA (Sacch. cerevisiae) IgA

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

ASCA (Sacch. cerevisiae) IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21



ASCA (Sacch. cerevisiae) IgM

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is a LDT and not covered by the accreditation of our laboratory.
as at: 11/22

Asialoglycoprotein rec. (ASGPR) IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is a LDT and not covered by the accreditation of our laboratory.
as at: 05/22

ASMA (GMA) IgG


sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

AT1A3 IgG

sample types: serum, EDTA, heparin, citrat plasma
alternative sample types: CSF
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is a LDT and not covered by the accreditation of our laboratory.
as at: 06/22

Basal ganglia IgG

sample types: serum, EDTA, heparin, citrat plasma
alternative sample types: CSF
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is a LDT and not covered by the accreditation of our laboratory.
as at: 05/22



Bile canaliculi IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is a LDT and not covered by the accreditation of our laboratory.
as at: 03/23

BP180 (NC16A-4x) IgA

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is a LDT and not covered by the accreditation of our laboratory.
as at: 11/22

BP180 (NC16A-4x) IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA, IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

BP230 IgA


sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is a LDT and not covered by the accreditation of our laboratory.
as at: 11/22

BP230 IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA, IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

C1q IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21



Calcium channels PQ-type IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: RIA
frequency: biweekly
reference value: see result report
as at: 12/21

cANCA IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Cardiac muscle IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Cardiac muscle intercalated discs IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Cardiolipin (AMA-M1) IgAGM

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Cardiolipin IgA

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21



Cardiolipin IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Cardiolipin IgM

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

CARPVIII IgG

sample types: serum, EDTA, heparin, citrat plasma
alternative sample types: CSF
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is a LDT and not covered by the accreditation of our laboratory.
as at: 06/22

CASPR1 IgG

sample types: serum, EDTA, heparin, citrat plasma
alternative sample types: CSF
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is a LDT and not covered by the accreditation of our laboratory.
as at: 06/22

CASPR2 IgG

sample types: serum, EDTA, heparin, citrat plasma
alternative sample types: CSF
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Cathepsin G IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21



CCP (cycl. citrull. peptide) IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Centrioles IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Centromeres IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Centromere protein B IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

CEP-1 IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

cN-1A (Mup44) IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21



Coeliac disease Profile IgA

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: EUROLINE
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Coeliac disease Profile IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: EUROLINE
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Coinlin; P80 (Few Nuclear Dots) IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Collagen type VII IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 04/22

Contactin1 IgG

sample types: serum, EDTA, heparin, citrat plasma
alternative sample types: CSF
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is a LDT and not covered by the accreditation of our laboratory.
as at: 06/22

CV2 (CRMP-5) IgG

sample types: serum, EDTA, heparin, citrat plasma
alternative sample types: CSF (LDT, not accredited)
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 05/22



CV2 IgG

sample types: serum, EDTA, heparin, citrat plasma
alternative sample types: CSF
sample volume: 1,5 ml
method: EUROLINE
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Cyclin I (PCNA) IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Cytoplasm HEp-2-cells IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Cytoplasm Profile 13 Ag IgG (M2, M2-3e, rib.P-Prot., Jo-1, SRP, PL-7, PL-12, EJ, OJ, Ha, Ks, Zo, Ro52)

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: EUROLINE
frequency: daily (mo-fri)
reference value: see result report
as at: 05/22

Cytoskeleton IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Desmoglein 1 IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA, IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21



Desmoglein 3 IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA, IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Desmosomes (intercellular substance) IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

DFS70 IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

DNA bound lactoferrin IgA


sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is a LDT and not covered by the accreditation of our laboratory.
as at: 05/22

DNA bound lactoferrin IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is a LDT and not covered by the accreditation of our laboratory.
as at: 08/23

Dopamin-receptors 2 IgG

sample types: serum, EDTA, heparin, citrat plasma
alternative sample types: CSF
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is a LDT and not covered by the accreditation of our laboratory.
as at: 06/22



DPPX IgG

sample types: serum, EDTA, heparin, citrat plasma
alternative sample types: CSF (LDT, not accredited)
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 05/22

dsDNA IgA (Crithidia luc.)

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is a LDT and not covered by the accreditation of our laboratory.
as at: 05/22

dsDNA IgAGM

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: RIA
frequency: biweekly
reference value: see result report
as at: 12/21

dsDNA IgG (Crithidia luc.)

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

dsDNA IgM (Crithidia luc.)

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is a LDT and not covered by the accreditation of our laboratory.
as at: 05/22

dsDNS-NcX IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21



EJ IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: EUROLINE
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Elastase IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

ENA Pool Plus IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

ENA Profile Plus 1 IgG (nRNP/Sm, Sm, SS-A, SS-B, Scl-70, Jo-1)


sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

ENA Profile Plus IgG EUROASSAY (nRNP/Sm, Sm, SS-A, SS-B, Scl-70, Jo-1)

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: EUROASSAY
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

ENA SLE Profile 2 (dsDNA, histones, nukleosomes, nRNP/Sm, Sm, SS-A, SS-B, Scl-70)

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21



Endomysium IgA

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Endomysium IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Enterocytes IgA

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is a LDT and not covered by the accreditation of our laboratory.
as at: 05/22

Enterocytes IgG


sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is a LDT and not covered by the accreditation of our laboratory.
as at: 05/22

Envoplakin IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Epidermal basement membrane IgA

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is a LDT and not covered by the accreditation of our laboratory.
as at: 11/22



Epidermal basement membrane IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

ERC1 IgG

sample types: serum, EDTA, heparin, citrat plasma
alternative sample types: CSF
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is a LDT and not covered by the accreditation of our laboratory.
as at: 06/22

Eye tissue IgAGM


sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is a LDT and not covered by the accreditation of our laboratory.
as at: 03/23

Flotillin 1/2 IgG

sample types: serum, EDTA, heparin, citrat plasma
alternative sample types: CSF
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is a LDT and not covered by the accreditation of our laboratory.
as at: 06/22

GABA-a-receptors IgG

sample types: serum, EDTA, heparin, citrat plasma
alternative sample types: CSF
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is a LDT and not covered by the accreditation of our laboratory.
as at: 06/22



GABA-b-receptors IgG

sample types: serum, EDTA, heparin, citrat plasma
alternative sample types: CSF
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

GAD 65 kDa IgG

sample types: serum, EDTA, heparin, citrat plasma
alternative sample types: CSF (LDT, not accredited)
sample volume: 1,5 ml
method: ELISA, IFA
frequency: ELISA biweekly, IFA daily (mo-fri)
reference value: see result report
as at: 05/22

GAD IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Ganglioside Profile 2 IgG (GM1, GM2, GM3, GD1a, GD1b, GT1b, GQ1b)


sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: EUROLINE
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Ganglioside Profile 2 IgM (GM1, GM2, GM3, GD1a, GD1b, GT1b, GQ1b)

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: EUROLINE
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

GBM (glomerular basement membrane) IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA, IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21



GFAP (Glial fibrillary acidic protein) IgG

sample types: serum, EDTA, heparin, citrat plasma
alternative sample types: CSF
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is a LDT and not covered by the accreditation of our laboratory.
as at: 05/22

Glia nuclear antibodies (AGNA) IgG

sample types: serum, EDTA, heparin, citrat plasma
alternative sample types: CSF (LDT, not accredited)
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 05/22

Gliadin-GAF-3X (Z-AGFA) IgA

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Gliadin-GAF-3X (Z-AGFA) IgG


sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

GluRD2 IgG

sample types: serum, EDTA, heparin, citrat plasma
alternative sample types: CSF
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is a LDT and not covered by the accreditation of our laboratory.
as at: 06/22

Glutamatreceptor (type AMPA) IgG

sample types: serum, EDTA, heparin, citrat plasma
alternative sample types: CSF (LDT, not accredited)
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 05/22



Glutamatreceptor (type NMDA) IgG

sample types: serum, EDTA, heparin, citrat plasma
alternative sample types: CSF
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Glycine receptors IgG

sample types: serum, EDTA, heparin, citrat plasma
alternative sample types: CSF
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is a LDT and not covered by the accreditation of our laboratory.
as at: 05/22

Goblet cells intestinal IgA

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Goblet cells intestinal IgG


sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Golgi apparatus IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Histones global IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21



Homer 3 IgG

sample types: serum, EDTA, heparin, citrat plasma
alternative sample types: CSF
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is a LDT and not covered by the accreditation of our laboratory.
as at: 06/22

Hu (ANNA-1) IgG

sample types: serum, EDTA, heparin, citrat plasma
alternative sample types: CSF (LDT, not accredited)
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 05/22

IA2 (Tyrosinphosphatase) IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: biweekly
reference value: see result report
as at: 12/21

ICA (islet cells) IgG


sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-thu)
reference value: see result report
as at: 12/21

IgLON5 IgG

sample types: serum, EDTA, heparin, citrat plasma
alternative sample types: CSF (LDT, not accredited)
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Insulin human IgG

sample types: serum sample volume: 1,5 ml
method: RIA
frequency: weekly
reference value: see result report
as at: 12/21



Intrinsic factor IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

ITPR1 IgG

sample types: serum, EDTA, heparin, citrat plasma
alternative sample types: CSF
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is a LDT and not covered by the accreditation of our laboratory.
as at: 06/22

Jo-1 IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

KCNA2 IgG


sample types: serum, EDTA, heparin, citrat plasma
alternative sample types: CSF
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is a LDT and not covered by the accreditation of our laboratory.
as at: 06/22

Kidney IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

KLHL11 IgG

sample types: serum, EDTA, heparin, citrat plasma
alternative sample types: CSF
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is a LDT and not covered by the accreditation of our laboratory.
as at: 11/22



Lacrinal gland IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is a LDT and not covered by the accreditation of our laboratory.
as at: 05/22

Laminin 5 (332) IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Laminin IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

LC-1 (Liver cytosol) IgG


sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Liver IgG IFA (globaltest, 6 biochips)

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Liver Profile (ALD) IgG (AMA-M2, M2-3e (BPO), Sp100, PML, gp210, LC-1, LKM-1, SLA/LP, SS-A, Ro-52, Scl-70, CENP A, CENP B, PGDH)

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: EUROLINE
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21



Liver Profile (ALD 9) IgG (AMA-M2, M2-3e (BPO), Sp100, PML, gp210, LKM-1, LC-1, F-Aktin Ro-52

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: EUROLINE
frequency: daily (mo-fri)
reference value: see result report
as at: 07/23

Liver Profile (AMA-M2, LKM-1, LC-1, SLA/LP)

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: EUROASSAY
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

LGI1 IgG

sample types: serum, EDTA, heparin, citrat plasma
alternative sample types: CSF (LDT, not accredited)
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 05/22

LKM-1 IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA, ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

LKM-2 IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is a LDT and not covered by the accreditation of our laboratory.
as at: 05/22

LMA (Liver cell membrane) IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is a LDT and not covered by the accreditation of our laboratory.
as at: 05/22



LRP 4 IgG

sample types: serum, EDTA, heparin, citrat plasma sample types 3: CFS
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is a LDT and not covered by the accreditation of our laboratory.
as at: 06/22

LSP (Liver-spec. proteine) IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Lysosomes IgG

sample types: serum sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/16

Ma (Ma1, Ma2/Ta) IgG

sample types: serum, EDTA, heparin, citrat plasma
alternative sample types: CSF (LDT)
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 05/22

MAG (Myelin assoz. Glykopr.) IgA

sample types: serum, EDTA, heparin, citrat plasma
alternative sample types: CSF
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is a LDT and not covered by the accreditation of our laboratory.
as at: 05/22

MAG (Myelin assoz. Glykopr.) IgG

sample types: serum, EDTA, heparin, citrat plasma
alternative sample types: CSF (LDT, not accredited)
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 05/22



MAG (Myelin assoz. Glykopr.) IgM

sample types: serum, EDTA, heparin, citrat plasma
alternative sample types: CSF (LDT, not accredited)
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 05/22

MBP (Myelobas. protein) IgG

sample types: serum, EDTA, heparin, citrat plasma
alternative sample types: CSF
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is a LDT and not covered by the accreditation of our laboratory.
as at: 05/22

MBP (Myelobas. protein) IgM


sample types: serum, EDTA, heparin, citrat plasma
alternative sample types: CSF
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is a LDT and not covered by the accreditation of our laboratory.
as at: 06/22

Melanocytes IgAGM

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is a LDT and not covered by the accreditation of our laboratory.
as at: 03/23

mGluR1 (metabotropic glutamate receptor 1) IgG

sample types: serum, EDTA, heparin, citrat plasma
alternative sample types: CSF
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is a LDT and not covered by the accreditation of our laboratory.
as at: 05/22



mGluR5 (metabotropic glutamate receptor 5) IgG

sample types: serum, EDTA, heparin, citrat plasma
alternative sample types: CSF
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is a LDT and not covered by the accreditation of our laboratory.
as at: 05/22

Mi-2 IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Mitosin (CENP-F) IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

MOG (Myelin-oligodendrocytes glycoprotein) IgG


sample types: serum, EDTA, heparin, citrat plasma
alternative sample types: CSF (LDT, not accredited)
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 05/22

MPO (Myeloperoxidase) IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

MSA-1 (NuMa-1) IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21



MSA-2 (HsEg5, spindle apparatus) IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

MSA-3 (CENP-F chromosomes ass. AG) IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Muscle striated IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

MuSK (musclespecific tyrosinkinase) IgG


sample types: serum sample volume: 1,5 ml
method: RIA
frequency: biweekly
reference value: see result report
as at: 12/21

MuSK IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFT
frequency: daily (Mo-Fr)
reference value: see result report
as at: 12/21

Muscarinic receptor (M2) IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is a LDT and not covered by the accreditation of our laboratory.
as at: 03/23



Muscarinic receptor (M3) IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is a LDT and not covered by the accreditation of our laboratory.
as at: 03/23

Myelin IgG

sample types: serum, EDTA, heparin, citrat plasma
alternative sample types: CSF (LDT, not accredited)
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 05/22

Myolemma IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is a LDT and not covered by the accreditation of our laboratory.
as at: 05/22

Myopathie Profile 4 IgG (Mi-2alpha/beta, TIF1gamma, MDA5, NXP2, SAE1, Ku, PM100, PM75, Jo-1, SRP, PL-7, PL12, EJ, OJ, Ro52, cN-1a, Ha, Ks, Zo)


sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: EUROLINE
frequency: daily (mo-fri)
reference value: see result report
as at: 05/22

Myosin IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Neuronal antigens IgG

sample types: serum, EDTA, heparin, citrat plasma
alternative sample types: CSF (LDT, not accredited)
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 05/22



Neurexin-3-alpha IgG

sample types: serum, EDTA, heparin, citrat plasma
alternative sample types: CSF
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is a LDT and not covered by the accreditation of our laboratory.
as at: 06/22

Neurochondrin IgG

sample types: serum, EDTA, heparin, citrat plasma
alternative sample types: CSF
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is a LDT and not covered by the accreditation of our laboratory.
as at: 06/22

Neurofascin 155 IgG

sample types: serum, EDTA, heparin, citrat plasma
alternative sample types: CSF
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is a LDT and not covered by the accreditation of our laboratory.
as at: 06/22

Neurofascin 186 IgG

sample types: serum, EDTA, heparin, citrat plasma
alternative sample types: CSF
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is a LDT and not covered by the accreditation of our laboratory.
as at: 06/22

Anti Neurofilaments IgG

sample types: serum, EDTA, heparin, citrat plasma
alternative sample types: CSF (LDT, not accredited)
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 06/22



Neuronale immunoblot IgG (amphiphysin, CV2, PNMA2 (Ma2/Ta), Ri, Yo, Hu, Recoverin, SOX1, Titin, Zic4, GAD65, Tr (DNER))

sample types: serum, EDTA, heparin, citrat plasma
alternative sample types: CSF
sample volume: 1,5 ml
method: EUROLINE
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

NOR (nucleus-organisator-region) 90 IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Nuclear membrane IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Nucleosomes IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

OJ IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: EUROLINE
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Ovary IgAGM

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21



Ovary teca cells IgAGM

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

pANCA IgA

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

pANCA IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Pancreas execution duct IgA

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is a LDT and not covered by the accreditation of our laboratory.
as at: 05/22

Pancreas execution duct IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is a LDT and not covered by the accreditation of our laboratory.
as at: 05/22

Pancreas acini IgA (CUZD1, GP2)

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21



Pancreas acini IgG (CUZD1, GP2)

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Parathyroid gland IgAGM

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is a LDT and not covered by the accreditation of our laboratory.
as at: 11/23

PCA (H/K-ATPase) IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

PCA (parietalcells) IgG


sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

PCA-2 (purkinje cell-cytopl.) IgG

sample types: serum, EDTA, heparin, citrat plasma
alternative sample types: CSF (LDT)
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 05/22

PDGH IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: EUROLINE
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21



Phosphatidyl-serin IgA

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Phosphatidyl-serin IgAGM

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Phosphatidyl-serin IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Phosphatidyl-serin IgM


sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Phospholipase A2-receptor IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA, ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Pituitary gland anterior lobe IgAGM

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is a LDT and not covered by the accreditation of our laboratory.
as at: 12/21



Pituitary gland posterior lobe IgAGM

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is a LDT and not covered by the accreditation of our laboratory.
as at: 12/21

PL-12 IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: EUROLINE
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

PL-7 IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: EUROLINE
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Plexus myentericus IgG


sample types: serum, EDTA, heparin, citrat plasma
alternative sample types: CSF (LDT, not accredited)
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 05/22

PM-1/PM-Scl IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Potassium channels

sample types: serum
CSF (LDT, not accredited) sample volume: 1,5 ml
method: RIA
frequency: weekly
reference value: see result report
as at: 05/22



PR3-hn-hr (proteinase3) IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Purkinje cell Profile (Yo, CDR2L, Tr, PRKCG, ARHGAP26, Homer-3, RGS8, RYR2, AP3B2) IgG

sample types: serum, EDTA, heparin, citrat plasma
alternative sample types: CSF (LDT, not accredited)
sample volume: 1,5 ml
method: EUROLINE
frequency: daily (mo-fri)
reference value: see result report
as at: 10/22

RANA (Rheumatoide arthritis nuclear antigen) IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is a LDT and not covered by the accreditation of our laboratory.
as at: 12/21

Recoverin IgG

sample types: serum, EDTA, heparin, citrat plasma
alternative sample types: CSF
sample volume: 1,5 ml
method: EUROLINE
frequency: daily (mo-fri)
reference value: see result report
as at: 11/22

Retina IgAGM

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is a LDT and not covered by the accreditation of our laboratory.
as at: 03/23

Rheumatoid factor IgA

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21



Rheumatoid factor IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Rheumatoid factor (class. Rheumatoid factor) IgM

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Rho-GTPase activating protein 26 IgG

sample types: serum, EDTA, heparin, citrat plasma
alternative sample types: CSF
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is a LDT and not covered by the accreditation of our laboratory.
as at: 06/22

Ri (ANNA-2) IgG

sample types: serum, EDTA, heparin, citrat plasma
alternative sample types: CSF (LDT, not accredited)
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 05/22

Ribosomal P-proteins IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 05/22

Ribosomes IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21



RNA-polymerase IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Ro-52 IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: EUROLINE
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Sa IgG

sample types: serum sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Sarcolemma IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is a LDT and not covered by the accreditation of our laboratory.
as at: 05/22

Scl-70 (DNA topoisomerase I) IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Signal recognition particle (SRP) IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: EUROLINE
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21



SLA/LP IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Sm IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

SOX-1 IgG

sample types: serum, EDTA, heparin, citrat plasma
alternative sample types: CSF
sample volume: 1,5 ml
method: EUROLINE
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Sp100, PML (nuclear dots) IgG


sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Salivary gland excretory ducts IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is a LDT and not covered by the accreditation of our laboratory.
as at: 11/23

Salivary gland acini IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 11/23



Salivary gland IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 11/23

Spermatozoa IgA

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Spermatozoa IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Spermatozoa IgM


sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

SS-A (Ro) 60 kDa

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

SS-B (La) IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21



ssDNS IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Steroid hormone prod. cells (ovary) IgAGM

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 05/22

Systemsclerosis IgG EUROLINE (Scl-70, CENP A, CENP B, RP11, RP155, Fibrillarin, NOR90, Th/To, PM-Scl100, PM-Scl75, Ku, PDGFR, Ro-52)

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: EUROLINE
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Testis IgAGM


sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is a LDT and not covered by the accreditation of our laboratory.
as at: 12/21

TG (thyroglobulin) IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA, IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 05/23

THSD7A IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
remarks:
as at: 12/21



Thymus IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is a LDT and not covered by the accreditation of our laboratory.
as at: 05/22

Thyroid microsomes IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Th/To IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: EUROLINE
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Titin IgG


sample types: serum, EDTA, heparin, citrat plasma
alternative sample types: CSF
sample volume: 1,5 ml
method: EUROLINE
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

To (7-2 RNP) IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: EUROLINE
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

TPO (thyreoperoxidase) IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21



Tr (DNER) IgG

sample types: serum, EDTA, heparin, citrat plasma
alternative sample types: CSF (LDT, not accredited)
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 05/22

TRAk (TSH-receptor) IgG

sample types: serum sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Transglutaminase IgA

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Transglutaminase IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Tropomyosin IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Tubular basement membrane IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21



U1-(n) RNP/Sm IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

U3-nRNP (Fibrillarin) IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA, EUROLINE
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Vascular endothelium IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is a LDT and not covered by the accreditation of our laboratory.
as at: 06/23

Vasopressin prod. cells IgAGM


sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is a LDT and not covered by the accreditation of our laboratory.
as at: 05/22

Vimentin IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Vinculin IgAGM

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is a LDT and not covered by the accreditation of our laboratory.
as at: 05/22



Yo (PCA-1) IgG

sample types: serum, EDTA, heparin, citrat plasma
alternative sample types: CSF
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

ZIC4 IgG

sample types: serum, EDTA, heparin, citrat plasma
alternative sample types: CSF
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is a LDT and not covered by the accreditation of our laboratory.
as at: 05/23

ZIC4 IgG

sample types: serum, EDTA, heparin, citrat plasma
alternative sample types: CSF
sample volume: 1,5 ml
method: EUROLINE
frequency: daily (mo-fri)
reference value: see result report
as at: 05/22

Zinc transporter (ZnT8)

sample types: serum
heparin, citrate plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

ZSCAN5B IgG

sample types: serum, EDTA, heparin, citrat plasma
alternative sample types: CSF
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is a LDT and not covered by the accreditation of our laboratory.
as at: 07/23



β2-glycoprotein IgA

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

β2-glycoprotein IgAGM


sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

β2-glycoprotein IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

β2-glycoprotein IgM

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21



Detection of

Circulating immune complexes C1q

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: weekly
reference value: see result report
as at: 12/21

CXCL13

sample types: CSF
sample volume: 1,5 ml
method: ELISA
frequency: weekly
reference value: see result report
as at: 01/19

Neurofilament (pNf-H) ELISA

sample types: serum CSF sample volume: 1,5 ml
method: ELISA
frequency: biweekly
reference value: see result report
remarks: Please note the instructions for preanalytics. The samples must be submitted frozen.
This analysis is not covered by the current accreditation scope.
as at: 09/24

Neurofilament (Nf-L) ELISA

sample types: CSF sample volume: 1,5 ml
method: ELISA
frequency: biweekly
reference value: see result report
remarks: Please note the instructions for preanalytics. The samples must be submitted frozen.
This analysis is not covered by the current accreditation scope.
as at: 09/24

pTau(181)

sample types: CSF
sample volume: 1,5 ml
method: ELISA
frequency: weekly
reference value: see result report
remarks: Please note the instructions for preanalytics. The samples must be submitted frozen.
This analysis is not covered by the current accreditation scope.
as at: 09/24

Total Tau

sample types: CSF
sample volume: 2,0 ml
method: ELISA
frequency: weekly
reference value: see result report
remarks: Please note the instructions for preanalytics. The samples must be submitted frozen.
This analysis is not covered by the current accreditation scope.
as at: 09/24



Detection of

β-amyloid (1-40) ELISA

sample types: CSF sample volume: 2,0 ml
method: ELISA
frequency: weekly
reference value: see result report
remarks: Please note the instructions for preanalytics. The samples must be submitted frozen.
This analysis is not covered by the current accreditation scope.
as at: 09/24

β-amyloid (1-42) ELISA

sample types: CSF sample volume: 2,0 ml
method: ELISA
frequency: weekly
reference value: see result report
remarks: Please note the instructions for preanalytics. The samples must be submitted frozen.
This analysis is not covered by the current accreditation scope.
as at: 09/24