



Specification List

Infectious Serology

General Information

The specifications list contains the range of analysis provided by the clinical immunological laboratory and informs you about the methods, sample types and volumes used, the frequency of analysis, reference values and, if necessary, preanalytical aspects. The specifications list is available on the internet at www.labor-stoecker.de.

Duration:

The samples are registered upon receipt in the laboratory. Preferably, analysis is performed or started on the same day to ensure that the report can be sent to the customer generally one day after receipt of the sample at the latest. The time period may be longer in individual cases if the analysis is not carried out daily or retesting of the sample or performing of a dilution series is required.

Reference range:

The reference values given in the specifications list are in accordance with the manufacturer's recommendations for the test system used and with current scientific and technological knowledge. However, reference ranges may change due to new scientific findings and technical developments in analytical techniques. Interpretation of results is always based on the reference ranges and additional remarks which are given on the report.

Measurement deviations:


The clinical immunological laboratory performs regular checks and maintenance to minimise variations and method-related deviations in the measurement results. Due to existing measurement deviations, borderline and weak-positive results or results that lie near to the reference value or other medical decision limits should be interpreted with care. Information about measurement deviations in individual quantitative analyses can be provided upon request. However, most analyses that are performed by the clinical immunological laboratory (antibody diagnostics) are considered qualitative.

Comparability of test results:

Results from different methods (e.g. immunofluorescence, ELISA, blot techniques) for detecting the same antibodies in a sample may differ. This can be due to the use of different antigens or to varying immobilisation of the same antigen to different carrier materials, which can influence the binding of the antibody to the antigen through conformational changes in the antigen or through steric inhibition.

Laboratory developed test (LDT):

The laboratory tests are usually CE marked. Tests that are not CE marked but validated by the laboratory are identified in the specification list as laboratory developed tests (LDT) and are not covered by the accreditation of the clinical immunological laboratory.



Antibodies against

Adenovirus IgA

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is not covered by the accreditation of our laboratory.
as at: 03/24

Adenovirus IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 03/22

Adenovirus IgM

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 03/22

Bartonella henselae IgG


sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Bartonella henselae IgM

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Bartonella quintana IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21



Antibodies against

Bartonella quintana IgM

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Bordetella parapertussis IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is a LDT and not covered by the accreditation of our laboratory.
as at: 11/23

Bordetella-FHA IgA

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Bordetella-FHA IgG


sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Bordetella-pertussis IgA

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: EUROLINE
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Bordetella-pertussis IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: EUROLINE
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21



Antibodies against

Bordetella-pertussis IgM

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Bordetella-pertussis-Toxin IgA

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Bordetella-pertussis-Toxin IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Borrelia afzelii IgG


sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 08/22

Borrelia afzelii IgM

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 08/22

Borrelia burgdorferi (USA) IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 08/22



Antibodies against

Borrelia burgdorferi (USA) IgM

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 08/22

Borrelia garinii IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: Westernblot
frequency: daily (mo-fri)
reference value: see result report
as at: 08/22

Borrelia garinii IgM

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: Westernblot
frequency: daily (mo-fri)
reference value: see result report
as at: 08/22

Borrelia IgM


sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA, EUROLINE/Westernblot
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Borrelia CSF/serum quotient IgG

sample types: serum and CSF
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Borrelia CSF/serum quotient IgM

sample types: serum and CSF
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21



Antibodies against

Borrelia plus VisE IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA, EUROLINE/Westernblot
frequency: daily (mo-fri)
reference value: see result report
as at: 08/22

Borrelia-RN-AT IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: EUROLINE
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Borrelia-RN-AT IgM

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: EUROLINE
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Brucella abortus IgG


sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Brucella abortus IgM

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Campylobacter jejuni IgA

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report remarks:
as at: 08/22



Antibodies against

Campylobacter jejuni IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA, ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Candida crusei IgAGM

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is a LDT and not covered by the accreditation of our laboratory.
as at: 05/22

Candida glabrata IgAGM

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is a LDT and not covered by the accreditation of our laboratory.
as at: 05/22

Candida parapsilosis IgAGM


sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is a LDT and not covered by the accreditation of our laboratory.
as at: 05/22

Candida tropicalis IgAGM

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is a LDT and not covered by the accreditation of our laboratory.
as at: 05/22

Chikungunya Virus IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA, IFA
frequency: biweekly
reference value: see result report
as at: 08/22



Antibodies against

Chikungunya Virus IgM

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA, IFA
frequency: biweekly
reference value: see result report
as at: 08/22

Chlamydia pneumoniae IgA

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA, MIF
frequency: daily (mo-fri)
reference value: see result report
as at: 08/22

Chlamydia pneumoniae IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA, MIF
frequency: daily (mo-fri)
reference value: see result report
as at: 08/22

Chlamydia pneumoniae IgM


sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA, MIF
frequency: daily (mo-fri)
reference value: see result report
as at: 08/22

Chlamydia psittaci IgA MIF

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: MIF
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Chlamydia psittaci IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: MIF
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21



Antibodies against

Chlamydia psittaci IgM

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: MIF
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Chlamydia trachomatis IgA

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA, MIF
frequency: daily (mo-fri)
reference value: see result report
as at: 08/22

Chlamydia trachomatis IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA, MIF
frequency: daily (mo-fri)
reference value: see result report
as at: 08/22

Chlamydia trachomatis IgM


sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA, MIF
frequency: daily (mo-fri)
reference value: see result report
as at: 08/22

Chlamydia -HP (pneumoniae, psittaci, trachomatis) IgA

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: EUROLINE/Westernblot
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is not covered by the accreditation of our laboratory.
as at: 08/22

Chlamydia-HP (pneumoniae, psittaci, trachomatis) IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: EUROLINE/Westernblot
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is not covered by the accreditation of our laboratory.
as at: 08/22



Antibodies against

CMV IgG avidity

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

CMV IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
remarks: This analysis is not intended to assess suitability for transfusion, transplantation or cell therapy.
as at: 05/22

CMV IgM

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
remarks: This analysis is not intended to assess suitability for transfusion, transplantation or cell therapy.
as at: 05/22

CMV CSF/serum quotient IgG


sample types: serum and CSF
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Coxsackie A16 IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 08/22

Coxsackie A16 IgM

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 08/22



Antibodies against

Coxsackie A7 IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 08/22

Coxsackie A7 IgM

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 08/22

Coxsackie B1 IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 08/22

Coxsackie B1 IgM

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 08/22

CXCL13 (direct)


sample types: CSF sample volume: 1,5 ml
method: ELISA
frequency: weekly
reference value: see result report
as at: 08/22

Dengue Virus 1-4 IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA, IFA
frequency: biweekly
reference value: see result report
as at: 12/21

Dengue Virus 1-4 IgM

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA, IFA
frequency: biweekly
reference value: see result report
as at: 12/21



Antibodies against

Dengue Virus NS1 (direct)

sample types: serum, EDTA, heparin plasma
sample volume: 1,5 ml
method: ELISA
frequency: biweekly
reference value: see result report
as at: 08/22

Diphtherie Toxoid IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

EBNA IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA, IFA
frequency: daily (mo-fri)
reference value: see result report
remarks: This analysis is not intended to assess suitability for transfusion, transplantation or cell therapy.
as at: 05/22

EBV sequence (EBV-CA IgG/IgM/avidity, EA IgG, EBNA)


sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
remarks: This analysis is not intended to assess suitability for transfusion, transplantation or cell therapy.
as at: 09/22

EBV-CA IgA

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
remarks: This analysis is not intended to assess suitability for transfusion, transplantation or cell therapy.
as at: 09/22

EBV-CA IgG Avidity

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA, IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21



Antibodies against

EBV-CA IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA, IFA
frequency: daily (mo-fri)
reference value: see result report
remarks: This analysis is not intended to assess suitability for transfusion, transplantation or cell therapy.
as at: 05/22

EBV-CA IgM

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA, IFA
frequency: daily (mo-fri)
reference value: see result report
remarks: This analysis is not intended to assess suitability for transfusion, transplantation or cell therapy.
as at: 05/22

EBV-CA CSF/serum quotient IgG


sample types: serum and CSF
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is not covered by the accreditation of our laboratory.
as at: 09/22

EBV-EA IgA

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
remarks: This analysis is not intended to assess suitability for transfusion, transplantation or cell therapy.
The analysis is not covered by the accreditation of our laboratory.
as at: 03/24

EBV-EA IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA, ELISA
frequency: daily (mo-fri)
reference value: see result report
remarks: This analysis is not intended to assess suitability for transfusion, transplantation or cell therapy.
as at: 05/22



Antibodies against

EBV-EA IgM

sample types: serum, EDTA plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
remarks: This analysis is not intended to assess suitability for transfusion, transplantation or cell therapy.
The analysis is not covered by the accreditation of our laboratory.
as at: 03/24

EBV-Profile 2 EUROLINE IgG (VCA gp125, VCA p19, EBNA-1, p22, EA-D)

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: EUROLINE
frequency: daily (mo-fri)
reference value: see result report
remarks: This analysis is not intended to assess suitability for transfusion, transplantation or cell therapy.
as at: 12/21

EBV-Profile 2 EUROLINE IgM (VCA gp125, VCA p19, EBNA-1, p22, EA-D)


sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: EUROLINE
frequency: daily (mo-fri)
reference value: see result report
remarks: This analysis is not intended to assess suitability for transfusion, transplantation or cell therapy.
as at: 12/21

Echinococcus granulosus IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA, EUROLINE/Westernblot
frequency: daily (mo-fri)
reference value: see result report
as at: 07/22

Enterovirus IgA

sample types: serum, EDTA, heparin plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is not covered by the accreditation of our laboratory.
as at: 08/22



Antibodies against

Enterovirus IgG

sample types: serum, EDTA, heparin plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is not covered by the accreditation of our laboratory.
as at: 08/22

Enterovirus IgM

sample types: serum, EDTA, heparin plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is not covered by the accreditation of our laboratory.
as at: 08/22

Hanta-Profile (HNTV, PUUV, DOBV) IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: EUROLINE
frequency: biweekly
reference value: see result report
as at: 09/22

Hanta-Profile (HNTV, PUUV, DOBV) IgM


sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: EUROLINE
frequency: biweekly
reference value: see result report
as at: 09/22

Hanta virus (HNTV, SNV, PUUV, DOBV, SEOV, SAAV) IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFT
frequency: biweekly
reference value: see result report
as at: 09/22

Hanta virus (HNTV, SNV, PUUV, DOBV, SEOV, SAAV) IgM

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFT
frequency: biweekly
reference value: see result report
as at: 09/22



Antibodies against

Hanta Virus Pool 1 IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: biweekly
reference value: see result report
as at: 12/21

Hanta Virus Pool 1 IgM

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: biweekly
reference value: see result report
as at: 12/21

Helicobacter pylori (CagA) IgA

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Helicobacter pylori IgA


sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA, EUROLINE/Westernblot
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Helicobacter pylori IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA, EUROLINE/Westernblot
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Hepatitis E IgA

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is not covered by the accreditation of our laboratory.
as at: 03/24



Antibodies against

Hepatitis E IgAGM

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is not covered by the accreditation of our laboratory.
as at: 03/24

Hepatitis E IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Hepatitis-E IgM

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

HHV-6 IgG


sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

HHV-6 IgM

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

HSV 1/2 Pool IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21



Antibodies against

HSV 1/2 Pool IgM

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

HSV 1/2 Pool CSF/serum quotient IgG

sample types: serum and CSF
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

HSV C1, G1+G2 IgG EUROLINE/WB

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: EUROLINE/Westernblot
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

HSV C1, G1+G2 IgM EUROLINE/WB


sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: EUROLINE/Westernblot
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

HSV-1 IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 03/23

HSV-1 IgM

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is a LDT and not covered by the accreditation of our laboratory.
as at: 03/23



Antibodies against

HSV-2 IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 03/23

HSV-2 IgM

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is a LDT and not covered by the accreditation of our laboratory.
as at: 03/23

Influenza A Virus IgA

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 09/22

Influenza A Virus IgG


sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 09/22

Influenza B Virus IgA

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 09/11

Influenza B Virus IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 09/22



Antibodies against

Japanese encephalitis virus IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: biweekly (Mo, Do)
reference value: see result report
as at: 12/21

Japanese encephalitis virus IgM

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: biweekly (Mo, Do)
reference value: see result report
as at: 12/21

Klebsiella pneumoniae IgA

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is a LDT and not covered by the accreditation of our laboratory.
as at: 05/22

Klebsiella pneumoniae IgG


sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is a LDT and not covered by the accreditation of our laboratory.
as at: 05/22

Klebsiella pneumoniae IgM

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is a LDT and not covered by the accreditation of our laboratory.
as at: 05/22

Leg. longbeachae (non-pneumophila) IgAGM

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 07/22



Antibodies against

Legionella pneumophila (serotype 1-7) IgAGM

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 07/22

Legionella pneumophila ST1-7 IgA

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Legionella pneumophila ST1-7 IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Legionella pneumophila ST1-7 IgM


sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Leishmania donovani IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Leishmania donovani IgM

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21



Antibodies against

Listeria monocytogenes 1/2a IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is not covered by the accreditation of our laboratory.
as at: 08/22

Listeria monocytogenes 1/2a IgM

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is not covered by the accreditation of our laboratory.
as at: 08/22

Listeria monocytogenes 4b IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is not covered by the accreditation of our laboratory.
as at: 08/22

Listeria monocytogenes 4b IgM


sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is not covered by the accreditation of our laboratory.
as at: 08/22

Measles Virus IgG avidity

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Measles Virus IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 09/22



Antibodies against

Measles Virus IgM

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 09/22

Measles Virus CSF/serum quotient IgG

sample types: serum and CSF
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Mumps Virus CSF/serum quotient IgG

sample types: serum and CSF
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Mumps Virus IgG


sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 09/22

Mumps Virus IgM

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 09/22

Mycoplasma hominis IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21



Antibodies against

Mycoplasma hominis IgM

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Mycoplasma pneumoniae IgA

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 06/22

Mycoplasma pneumoniae IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 06/22

Mycoplasma pneumoniae IgM


sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 06/22

Parainfluenza Virus type 1-4 Pool IgA

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Parainfluenza Virus type 1-4 Pool IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21



Antibodies against

Parvovirus B19 IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA, EUROLINE
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Parvovirus B19 IgM

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA, EUROLINE
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Plasmodium IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: biweekly
reference value: see result report
remarks: This analysis is not intended to assess suitability for transfusion, transplantation or cell therapy.
as at: 05/22

RSV IgA


sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 09/22

RSV IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 09/22

RSV IgM

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is not covered by the accreditation of our laboratory.
as at: 03/24



Antibodies against

Rubella Virus IgG avidity

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 09/22

Rubella Virus IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 09/22

Rubella Virus GP IgM

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Rubella Virus CSF/serum quotient IgG


sample types: serum and CSF
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

SARS-CoV-2-(spike-protein) QuantiVac IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
alternative sample types: dried blood spots (not covered by the accreditation of our laboratory)
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 09/22

SARS-CoV-2-NCP IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
alternative sample types: dried blood spots (not covered by the accreditation of our laboratory)
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21



Antibodies against

SARS-CoV-2-NCP IgM

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is not covered by the accreditation of our laboratory.
as at: 05/22

SARS-CoV-2-Neutralisation assay

sample types: serum, EDTA, heparin plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

SARS-CoV-2-Spike-Protein IgA

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 09/22

Schistosoma mansoni IgG


sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA, ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Schistosoma mansoni IgM

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA, ELISA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is not covered by the accreditation of our laboratory.
as at: 08/22

TBE Virus CSF/serum quotient IgG

sample types: serum and CSF
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21



Antibodies against

TBE virus CSF/serum quotient IgM

sample types: serum and CSF
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

TBE Virus IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA, IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

TBE Virus IgM

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA, IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Toxoplasma gondii IgA


sample types: serum, EDTA plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
remarks: This analysis is not intended to assess suitability for transfusion, transplantation or cell therapy.
as at: 12/21

Toxoplasma gondii IgG avidity

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 08/22

Toxoplasma gondii IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA, IFT
frequency: daily (mo-fri)
reference value: see result report
remarks: This analysis is not intended to assess suitability for transfusion, transplantation or cell therapy.
as at: 08/22



Antibodies against

Toxoplasma gondii IgM

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA, IFT
frequency: daily (mo-fri)
reference value: see result report
remarks: This analysis is not intended to assess suitability for transfusion, transplantation or cell therapy.
as at: 08/22

Toxoplasma gondii CSF/serum quotient IgG

sample types: serum and CSF
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Treponema pallidum IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA, IFA, EUROLINE
frequency: daily (mo-fri)
reference value: see result report
remarks: This analysis is not intended to assess suitability for transfusion, transplantation or cell therapy.
as at: 05/22

Treponema pallidum IgGM


sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
remarks: This analysis is not intended to assess suitability for transfusion, transplantation or cell therapy.
as at: 05/22

Treponema pallidum IgM

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA, IFA, EUROLINE
frequency: daily (mo-fri)
reference value: see result report
remarks: This analysis is not intended to assess suitability for transfusion, transplantation or cell therapy.
as at: 05/22

Treponema pallidum CSF/serum quotient IgG

sample types: serum and CSF
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21



Antibodies against

Ureaplasma urealyticum IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

Ureaplasma urealyticum IgM

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

VZV IgA

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 09/22

VZV IgG avidity


sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 09/22

VZV IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA, IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

VZV IgM

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA, IFA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21



Antibodies against

VZV CSF/serum quotient IgG

sample types: serum and CSF
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
as at: 12/21

West Nile Virus IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: biweekly
reference value: see result report
as at: 12/21

West Nile Virus IgM

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: biweekly
reference value: see result report
as at: 12/21

West-Nil-Virus IgG


sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is not covered by the accreditation of our laboratory.
as at: 09/22

West-Nil-Virus IgM

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: daily (mo-fri)
reference value: see result report
remarks: The analysis is not covered by the accreditation of our laboratory.
as at: 09/22

Yellow fever virus IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: biweekly
reference value: see result report
remarks: The analysis is not covered by the accreditation of our laboratory.
as at: 05/22



Antibodies against

Yellow fever virus IgM

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: biweekly
reference value: see result report
remarks: The analysis is not covered by the accreditation of our laboratory.
as at: 05/22

Yersinia enterocolitica IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA, Westernblot
frequency: daily (mo-fri)
reference value: see result report
as at: 08/22

Zika Virus IgG

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: IFA
frequency: biweekly
reference value: see result report
remarks: The analysis is not covered by the accreditation of our laboratory.
as at: 05/22

Zika Virus IgM


sample types: serum, EDTA-, Heparin-, Citrat-Plasma
sample volume: 1,5 ml
method: IFA
frequency: biweekly
reference value: see result report
remarks: The analysis is not covered by the accreditation of our laboratory.
as at: 05/22

Zika-virus IgA

sample types: serum, EDTA, heparin plasma
sample volume: 1,5 ml
method: ELISA
frequency: biweekly
reference value: see result report
as at: 09/22

Zika-virus IgAM

sample types: serum, EDTA, heparin, citrat plasma
sample volume: 1,5 ml
method: ELISA
frequency: biweekly
reference value: see result report
as at: 09/22



Antibodies against

Zika-virus IgG

sample types: serum, EDTA, heparin plasma
sample volume: 1,5 ml
method: ELISA
frequency: biweekly
reference value: see result report
as at: 09/22

Zika-virus IgM

sample types: serum, EDTA, heparin plasma
sample volume: 1,5 ml
method: ELISA
frequency: biweekly
reference value: see result report
as at: 09/22